

Title: Recovery from stunting and cognitive outcomes in young children: Evidence from the South African Birth to Twenty Cohort Study.

Short title: Recovery from stunting and cognitive outcomes

Authors: Daniela Casale and Chris Desmond

(Forthcoming in 2015 special issue of *Journal of Developmental Origins of Health and Disease*)

Abstract

In this study we analyse the implications for cognitive function of recovery from stunting in early childhood. More specifically, we test whether children who met the definition for stunted at age two, but not at age five, perform better in cognitive tests than children who remain stunted over this period. The sample is drawn from the Birth to Twenty Cohort Study, a prospective dataset of children born in 1990 in urban South Africa. The measure of cognitive function that we use is based on the Revised-Denver Prescreening Developmental Questionnaire implemented when the children were age five. We employ multivariate regression in the analysis to control for child-specific characteristics, socio-economic status, the home environment, and caregiver inputs. We find that recovery from stunting is not uncommon among young children in our sample. However, children who recover from stunting by age five still perform significantly worse on cognitive tests than children who do not experience early malnutrition, and almost as poorly as children who remain stunted. These findings suggest that the timing of nutritional inputs in the early years is key in a child's cognitive development, with implications for school readiness and achievement.

Keywords: recovery from stunting; preschool cognitive function; birth cohort; South Africa

1. Introduction

Chronic child malnutrition remains a pervasive problem, with long-term implications for affected children. Stunting, measured as height-for-age Z-score (HAZ) of more than two standard deviations below the median HAZ of the reference population, is a useful marker of chronic malnutrition. Globally, an estimated 171 million children under 5 years of age were stunted in 2010.¹ Stunting early in life has been associated with a range of negative long-term outcomes, including impaired cognitive development, poor school outcomes, reduced earnings in adulthood and poor maternal reproductive health outcomes.²⁻⁴

The association between early life stunting and long-term outcomes is well-established. Where debate continues is in regards to the importance of timing (within the early years) and the persistence of stunting. Children in the poorer regions of the world are on average born shorter and grow more slowly, leading to a peak in stunting at 24 months, after which stunting prevalence either remains stable or begins to decline.⁵ The extent to which the children who are stunted at 24 months can catch up from this poor start and achieve similar adult heights as the reference population continues to be debated.⁶⁻¹⁰ It has been argued that if children can catch up then a narrow policy focus on what is now commonly referred to as the first 1,000 days (pregnancy and the first two years of life) is not warranted and should be broadened to the entire preschool period.⁸ However, even if complete catch up is possible (and normal adult height attained) and it is shown that later life interventions support such catch up, the question of whether later growth can undo the harm caused by stunting within the first 1,000 days remains.

The debate centers on the importance of the first 1,000 days for brain development. During pregnancy the process of neurulation leads to the development of the basic architecture of the brain. In early childhood the areas associated with vision and hearing, then language and

speech, and eventually higher cognitive function, undergo rapid synaptogenesis.¹¹ These foundational processes are energy intensive, making brain development highly susceptible to the negative consequences associated with malnutrition (both pre and postnatal).¹² As the development of the brain is cumulative, early difficulties can have life-long implications.¹³ A number of researchers have sought to identify if there are critical phases from a nutrition perspective within the first 1,000 days. Glewwe and King¹⁴, for example, investigate the impact of the timing of poor nutrition within the first two years of life on cognitive development and conclude that poor nutrition in the period 18-24 months had the most significant consequences. The results in Hodinott et al from Guatemala suggest that nutritional supplementation during 0-36 months of age, but not 36-72 months of age, resulted in higher average wages among adult men.¹⁵

Given that the period of rapid brain development continues throughout the early years, a small but growing literature has examined the extent to which catch-up growth after the first 1,000 days, measured as a recovery from stunting, mitigates the impact of stunting on cognitive development. Mendez and Adair¹⁶ using cohort data from Cebu in the Philippines found that children who catch up between 2 years and follow up at 8 and 11 years do worse at school than children who were never stunted, although less so than those who remain stunted. Crookston et al.¹⁷, using data from four developing countries, found a similar pattern; catch up between 1 year and 8 years appeared to reduce, but not eliminate, the deficit associated with early stunting. However, Crookston et al.^{18, 19} using data from Peru report that children who catch up between 6-18 months and 4.5-6 years do the same in cognitive tests as children who were never stunted, and better than children who remain stunted. The inconsistencies in reported results highlight the need for further analysis.

Using data from a South African longitudinal cohort, the Birth to Twenty Study, we examine whether recovery from stunting influences cognitive function. In previous work we found no association between stunting at 2 years and social maturity at 4 years but a large and significant association between stunting at 2 years and cognitive function at 5 years.²⁰ In this paper, we compare cognitive function among children who were stunted at 2 years but not at 5 years, with children who were stunted in both periods and children who were not stunted in either period.

The Birth to Twenty Study (Bt20) in South Africa is well-suited to examine the impact of recovery from stunting. Bt20 has both high rates of stunting and high rates of recovery from stunting.⁹ Moreover an extensive assessment of the child's cognitive development is conducted prior to school enrollment. There has been little prior research on stunting and its links to other aspects of child development in preschool-aged children using large sample data. The focus has been on schooling outcomes or cognitive tests conducted with school-going children. However, examining children once in the schooling system is complicated by various school-related confounders, such as school quality, and by the possibility that parents of smaller children delay enrollment.

Stunting and cognitive development may be correlated not only because of the impact of the former on the latter, but because they are both influenced by the environment (especially the home environment in early childhood) and caregiver inputs. A further advantage of Bt20 is that a wealth of data was collected on maternal/caregiver and household variables which we are able to use to account for these confounding factors in multivariate analysis.

2. Methods

Data and sample

The data are drawn from a birth cohort study from South Africa's largest metropolitan area. Referred to as Birth to Twenty (Bt20), this longitudinal study covered children born over a 7-week period between April and June of 1990 in private and public hospitals or clinics in the greater Johannesburg-Soweto area. The full sample comprises 3273 singleton births, however at any one interview point data is collected from roughly 1600 to 2200 participants. The attrition rate is amongst the lowest recorded for birth cohort studies in developing countries.^{21, 22}

Norris et al.²¹ describe attrition in the Bt20 study in detail. The main reason for participants being lost to follow-up appears to be child/family mobility, with child and caregiver mortality accounting for a very low share of the attrition. White South Africans, who are more likely to live in wealthier, suburban areas of the city, and who would have been more likely to give birth in private hospitals, are under-represented in the sample due to low enrolment in the study and subsequent non-response.^{22, 23} The Bt20 sample in follow-up waves therefore is most representative of children from the black African population group who were born in, and remained residents of, Johannesburg over the period.

Information was collected from mothers at antenatal clinics, at delivery centres, and thereafter generally once a year through face-to-face interviews with the caregiver and child. In our analysis, we use data from the delivery reports, year 2 (n=1839), year 4 (n=1858) and year 5 (n=1586). The number of children for whom there is non-missing data on our key explanatory variables, namely height-for-age at 2 years and 5 years, is 1574. However, missing data on the outcome variable and the control variables in the multivariate analysis result in a regression sample of between 1019 and 666, depending on which controls are included. We discuss the

implications of a diminishing sample for the mean values of key variables and the regression estimations in the Results section below.

Outcome measure

Our measure of cognitive function is based on the Revised-Denver Prescreening Developmental Questionnaire (R-DPDQ).²⁴ The assessment in Bt20 comprised 32 items covering the child's personal-social, fine motor, gross motor, language and cognitive abilities at age five. During the assessment some questions were asked of the caregiver, but the largest part of the questionnaire involved tests conducted by the interviewer. Some items were modified to be culturally appropriate (for example, the word 'hedge' was replaced with 'fence').²⁵ A full list of items is shown in Table 1 (for the detailed explanation of each question/test in the questionnaire, consult Hsiao and Richter²⁵).

[Table 1 about here]

Earlier research found a significant correlation between the Denver Development Screening Test and the Griffiths Mental Development Scales among black African preschool children in South Africa.²⁶ More recently, using Bt20 data, Hsiao and Richter²⁵ report strongly significant associations between performance in the R-DPDQ at age five and various other outcomes, namely scores on the Raven's Coloured Progressive Matrices and the Connor's Teacher Rating Questionnaire at age seven, age of first entry to school, and grade repetition by Grade 6. Internal consistency for the R-DPDQ measure in the Bt20 sample is 0.71, measured by Cronbach's α .

An overall score was calculated by adjusting the total raw score by the child's chronological age, and replacing missing values on individual items by the series mean. The majority of children had no missing data on the 32 items (73 per cent), and for those who did,

most of them were missing data on only one item, with the maximum number of missing items per individual being three. The mean value for the R-DPDQ at age 5 is 43.74 (SD=4.68, n=1232).

Key explanatory variable

The key variables of interest are stunting status at ages two and five, with stunting indicated by a height-for-age z-score (HAZ) of < -2 SD from the mean of the reference population, using WHO Child Growth Standards.²⁷ We use stunting at age two as the first point in the analysis as research has shown that the prevalence of stunting peaks at around 24 months^{5, 8}; the second point in the analysis is age five where we have information on both anthropometry and cognitive development. For our sample of children who had non-missing data on HAZ at ages two and five (n=1574), mean HAZ in 1992 when the children were aged two is -1.135 (SD=1.084) and mean HAZ in 1995 when the children were aged five is -0.664 (SD=0.895). The prevalence of stunting in this sample of children is 19.44 per cent at age two and 6.04 per cent at age five.

This pattern of rapid increases in stunting until 2 years followed by substantial recovery has been observed in other South African data. The National Health and Nutrition Examination Survey (2012), for example, found prevalence rates of 25.9 for girls and 26.9 for boys among 0-3 year olds, but 9.5 and 13.5 respectively among 4-6 year olds.²⁸ Although prevalence rates in the Bt20 sample are somewhat lower than the national figures cited here, this would be expected as the prevalence of stunting in urban areas has been found to lower than in rural areas in South Africa.²⁹

To estimate whether recovery from stunting between ages two and five resulted in better performance on cognitive test scores, we divide our sample of children into four groups

following the nomenclature of Mendez and Adair¹⁶: 1) not stunted at 2 y and not stunted at 5 y - labelled “neither”; 2) stunted at 2 y and 5 y - “persistent”; 3) not stunted at 2 y but stunted at 5 y - “late incident”; and 4) stunted at 2 y but not stunted at 5 y –“catch-up”. There is some debate as to what constitutes catch up in height in young children and whether or not it should be defined in relation to an external reference group^{6, 30} For simplicity, and for the sake of comparability with other studies, we use the definition most commonly used in the literature on growth and cognitive function, namely a recovery from stunting.^{16, 18, 31} However, in the final section of this paper we discuss the possibility of using alternative definitions.

Data analysis and control variables

Linear regression was used to estimate the relationship between stunting status in early childhood and the scores on the cognitive development test at 5y. The main independent variables of interest are introduced as dummy variables, “persistent” stunting at 2y and 5y, “late incident” stunting at 5y, and “catch-up” or stunting at 2y but not at 5y, where “neither” stunted at 2y or 5y is the omitted category.

Stunting is likely to be correlated with a number of other factors which may also be related to cognitive function. We therefore include an extensive set of controls in the regressions to try to minimise confounding effects. In addition to the unadjusted model, we show the results from models adjusted sequentially for groups of variables representing ‘characteristics at birth’, ‘socio-economic status’, and ‘the home environment/caregiver inputs’.

Specifically, characteristics at birth include the sex of the child and the child’s birth weight. Socio-economic status is captured by an asset score (based on six items in the survey - fridge, car, washing machine, television, phone and radio), maternal age (years) and maternal education (years of schooling). A dummy variable for the black African population group is

included, where the omitted category consists of Coloureds, Indians and Whites. Decades of institutionalised discrimination under *apartheid*, which affected black Africans most severely, resulted in racial disparities in almost all aspects of socio-economic life in South Africa, and the aforementioned socio-economic status variables may not capture this inequality adequately.

As discussed in the introduction key confounders not frequently adjusted for in such analyses are the household environment and the caregiver's inputs in the child's development, referred to in the economics literature as a 'preference for child quality'.¹⁴ We have a variety of variables in our very rich dataset which we can use to try to account for these. We include whether the mother was the main caregiver, birth order, birth spacing (indicating a birth within 24 months of the index child), how much time the mother/caregiver spent playing with the child each day, whether the mother/caregiver was trying to teach the child anything at the time, how often the father (or other man important to the child) spent playing with the child, and whether the child had any "playthings, bought toys or things (the caregiver) has made or given him/her to play with".

The variables described above (except for those captured at birth), were from the year 2 data round. In a final regression, we include two additional variables from year 4 to capture a change in the environment, namely whether the asset score (based on the same six items in year 2) showed no change or whether it increased between 2y and 4y (where the omitted category represents a decrease in assets).

A number of alternative measures of socio-economic status and the home environment were tested for significance in the cognitive function equation, among them, crowding in the household (the number of people per sleeping room), mother's height, paternal education, household income quintiles, maternal depression at 6 months, and an index of the quality of the

mother/child relationship. This latter variable was based on six items recorded by the interviewer, namely whether the child appeared clean and well looked after; seemed happy; appeared confident and secure in the mother's presence; and whether the mother seemed unhappy and worn down by worries and troubles; demonstrated any negative feelings towards the child; appeared to be confident and assured in her care and management of child; and showed affection towards the child.

None of these variables were found to be independently associated with cognitive function in our sample after controlling for the other factors described above, so they were not included in the final regressions for the sake of parsimony. (Details of these additional tests are available in Casale, Desmond and Richter²⁰). The mean values of the covariates included in the final analysis are shown in Table 2.

[Table 2 about here]

3. Results

Table 3 presents stunting status and mean height-for-age z-scores (HAZ) at 2y and 5y. The frame on the left-hand side of the table shows that at 2y, the 19.4 per cent of children who were stunted had a mean HAZ of -2.72 , and the 80.6 per cent who were not stunted had a mean HAZ of -0.751 . The right-hand side frame shows stunting status and mean HAZ at 5y for these same children. Of the full sample of children, 79.3 per cent were not stunted at 2y or 5y ("neither") and mean HAZ at 5y for this group is -0.425 . Another 4.7 per cent were stunted at both 2y and 5y ("persistent") with mean HAZ at 5y of -2.464 . Only a small percentage of children, 1.3 per cent, exhibit late incident stunting and mean HAZ at 5y for this group is -2.216 . The remaining 14.7 per cent constitute those who were stunted at 2y but not at 5y. The mean

HAZ at 5y for this “catch-up” group is -1.287, which is just less than a SD below the mean for those who were not stunted in either year, but over one SD higher than the mean of those who were stunted in both years or just in year 5.

[Table 3 about here]

While late incident stunting is not common in this sample, the rate of recovery from stunting is high. About 75 per cent of children who were stunted at 2y had a HAZ score of ≥ -2 SD at 5y. Furthermore, this high rate of recovery does not appear to be an artefact of small changes around the cut-off. The mean change in HAZ between 2y and 5y for the catch-up group is 1.343 (SD=0.864), none had a change in HAZ of less than 0.2 SD, and only 21 of the 231 children or 9.1 per cent had a change in HAZ of between 0.2 SD and 0.5 SD.

For the late incident group, however, the mean change in HAZ between 2y and 5y is -0.532 (SD=0.225). Among the 20 late incident cases only 1 had a change in HAZ of less than -0.2 SD but another 8 out of 20 (or 40 per cent) had a change in HAZ of between -0.2 and -0.5 SD. While this still suggests a considerable fall off in growth, we are cautious about the results for this group given the low sample size and the smaller changes around the cut-off.

Table 4 displays the unadjusted regression estimates, as well as the estimates from multivariate linear regressions where an increasing number of covariates is introduced with each model. Here, we restrict the analysis to the final regression sample of $n=666$ so that we can observe the varying effects of the covariate sets on the results.

[Table 4 about here]

The unadjusted estimates on the final regression sample suggest that children who are persistently stunted score 3.39 points less on the cognitive measure than children who were neither stunted at 2y or 5y, and this is significant at the 1 per cent level. Children with late

incident stunting also score less, but by a very small amount (0.36 points), and the coefficient is not significant. Those who recovered from stunting do considerably worse; they score 2.26 points less than children who were neither stunted at 2y or 5y, and this result is strongly significant at the 1 per cent level.

As we would expect, controlling for an increasing number of covariates reduces the coefficients on the stunting status variables, with birth characteristics and socio-economic status having the largest effects. Overall, the coefficient for children with persistent stunting drops from -3.39 to -2.51, and the coefficient for the catch-up group drops from -2.26 to -1.61, but both remain strongly significant. The coefficient on late incident stunting remains small and statistically insignificant.

Although the different size effects for the catch-up and persistent stunting groups suggest that the catch-up group do not perform as poorly as the persistently stunted group, the coefficients are not significantly different from each other ($F(1, 644) = 1.16$; $\text{Prob} > F = 0.283$). We also tested whether the inclusion of a variable capturing severe stunting ($\text{HAZ} < -3 \text{ SD}$) at 2y affected the results. We might expect this if, for example, children who catch up were not as badly stunted to begin with (or similarly, if those who are persistently stunted were more severely stunted at 2y). Using Model V controls, we find that the coefficient on severe stunting at 2y is not significantly correlated with cognitive function at 5y, and the coefficients on “persistent” and “catch-up” remain significant and largely unchanged at -2.61 and -1.65 respectively (compared to -2.51 and -1.61 without controlling for the severity of stunting).

A number of the controls are significantly related to cognitive function. Girl children do better than boy children and birth weight has an independent positive effect on cognitive function. Of the set of socio-economic status variables, the asset index and the dummy variable

for black African children are significant, with positive and negative effects respectively. Mother's education had a significant effect in Model III, but in Model IV when variables representing the home environment and caregiver inputs are included, the coefficient is no longer significant. This suggests that the positive effect of mother's education on cognitive function operates through some of these other variables.

Of the home environment/caregiver variables, whether the mother is the main caregiver is important, as is whether the mother/caregiver played with the child for at least an hour a day. Whether there were toys bought or made for the child in the home also has a strong positive effect on cognitive development. Birth spacing, however, has a negative effect on cognitive function, suggesting that another birth within 24 months of the index child places pressure on parental and other resources. A change in the asset index between 2y and 4y had no impact on the child's cognitive score.

As noted above, our sample size diminishes quite substantially from 1574 to 1019, due to missing values on the cognitive function score, and from 1019 to 666, with the inclusion of the covariates. To investigate the implications of this loss, we summarise in Table 5 the mean values of some the key variables for the three different samples. The most substantial change is in the proportion of Black children in the sample, which increases from 76 per cent to 89 per cent. There is very little difference across the samples in, for example, mean birth weight, mother's age, and the cognitive score itself.

The prevalence of stunting at 2y does increase by 3 percentage points with the loss of cases, indicating that taller children are being lost from the analytical sample. However, the loss of taller children does not appear to change the overall findings of the research. We attempt to show this in Table 6 by comparing the results on the unadjusted regression for the final

regression sample of 666 children (Model I in Table 4) with the results of an unadjusted regression on the sample of 1019 children for which we have information on cognitive function. While the size effects are larger for the persistent stunting variable as cases are dropped, there is very little difference in the coefficients on the ‘catch-up’ variable (-2.192 vs -2.258). Importantly, the substantive findings are consistent; children with persistent stunting do worse than those who were not stunted, as do the children who recovered from stunting between 2y and 5y. This gives us confidence that the results are not being driven by the non-random exclusion of cases due to the introduction of the covariates.

[Tables 5 and 6 about here]

4. Discussion

In our sample of children from the Bt20 cohort study we find that 75% of children who were stunted at 2y were no longer stunted at 5y. The children who recovered from stunting at 2y still performed significantly worse on our cognitive assessment at 5y than did children who were never stunted. This relationship remains significant when we include a full set of controls for potential confounding variables. Children who are no longer classified as stunted perform better than children who remain stunted, but the difference is not significant. Our results appear to provide further evidence that the negative implications for cognitive development of growth retardation within the first 1,000 days are not mitigated by subsequent catch-up growth.

Our result that children who recover from early stunting perform worse on cognitive assessments than those who were never stunted is similar to that of Mendez and Adair¹⁶ and Crookston et al’s¹⁷ multi-country analysis. However, in both of these studies the better performance of the catch-up group in relation to the persistently stunted group was significant. In our study we find the same pattern, but the difference is not significant. In contrast to these

findings, Crookston et al.^{18,19} found that children from a Peruvian cohort who experienced catch-up growth had cognitive scores similar to children who were never stunted. The Crookston et al multi-country study includes analysis of the same Peruvian Young Lives dataset, although in the multi-country analysis they focus on outcomes at age 8, rather than at age 4.5 – 6 years as was the case in the single country study.

Direct comparison to either the multi-country or single country Crookston et al analyses is complicated by the age at which early stunting was assessed. Unlike in our and the Mendez and Adair¹⁶ analysis, in which stunting was assessed at 2 years, the Young Lives data used by Crookston et al.¹⁷⁻¹⁹ includes stunting measured at 6-18 months.. The finding by Glewwe and King¹⁴ that the 18-24 month period is critical suggests that the differences in timing may lead to differences in the assessment of the importance of catch-up growth. Some proportion of the children who were assigned to the catch-up group in the Crookston et al.¹⁷⁻¹⁹ analysis may have started to recover from stunting in the second year of life. Had these children been assessed at 2y they may not have been classified as stunted. The inclusion of children who experience early catch up may be pulling the average cognitive scores for the overall catch-up group upward. Similarly, the children who become stunted between 18 and 24 months may be pulling the average score for never stunted children downward.

Although our results on stunting recovery are similar to Mendez and Adair,¹⁶ we do not find, as they did, that severe stunting played an independent role in predicting cognitive scores. Berkman et al.³¹ also report a significant impact of severe stunting in the second year of life on cognitive development. Our failure to find an independent effect of severe stunting may relate to the differences in age of children at follow-up. We examined cognitive development at 5 years of age, before the start of formal schooling. Mendez and Adair¹⁶ have information on cognitive

function at 8 and 11 years and Berkman et al.³¹ at 9 years of age. A supportive schooling environment (and continued growth in height) may help mitigate the impact of stunting and children who were moderately stunted may be more likely to benefit. Notably, the importance of severity increased in the Mendez and Adair¹⁶ analysis between 8 and 11 years.

To allow for comparison with other work in this field we refer to children who were stunted at 2y but not at 5y as having experienced catch-up growth. There are alternative definitions of catch-up growth. Cameron, Preece and Cole,⁶ for example, define catch up as an improvement in HAZ greater than that which would be predicted by regression to the mean, operationalized in the conditional growth measures used in, for example, Adair et al.³² Boersma and Wit³³ distinguish between three types of catch-up growth: type A where a child experiences an increase in height velocity such that their height deficit is swiftly eliminated; type B where the period of growth is prolonged; and type C which is a combination of A and B. Moreover, they draw a distinction between catch-up growth and the resumption of a normal growth velocity.

Using the Cameron, Preece and Cole⁶ definition of catch-up growth or identifying those children who could be classified as having type A catch-up, may well alter our findings on the association of catch-up growth and later cognitive outcomes. Recovery from stunting is a weak definition of catch up. It is possible that a child who is stunted at 2y and then resumes a normal growth velocity will no longer be defined as stunted at 5y. As children get older (and taller) a standard deviation in the reference population is associated with a larger and larger absolute difference in height. A child can remain the same absolute amount shorter than the mean and recover from stunting because relative to the mean that absolute difference is no longer 2 standard deviations below. It is possible that they even fall further behind in absolute height and still recover from stunting, because the change in the size of a standard deviation with age is

large enough.^{34, 35} Our use of the most common definition of catch up has allowed for direct comparison, but further work is clearly needed to investigate the implications of different definitions of catch up on the conclusion that it does not lead to improved cognition.

Our paper provides further evidence of the importance for cognitive development of linear growth, and by implication, nutrition, in the first two years of life. However, additional research is required to examine if the age at which early stunting is measured influences the assessment of the impact of catch-up growth on subsequent cognitive outcomes. Moreover, further research is required to investigate the extent to which the negative cognitive impact fades over time, and if it fades equally for moderately and severely stunted children. Finally, alternative definitions of catch up need to be tested, before it is concluded that catch-up growth does not mitigate the impact of early growth retardation on later cognitive outcomes.

Acknowledgements

We would like to thank Celia Hsiao, Shane Norris and Linda Richter for their helpful suggestions and guidance during the preparation of this paper.

Financial support

Birth to Twenty is funded by the Wellcome Trust, the University of the Witwatersrand, the South African Medical Research Council and the South African Human Sciences Research Council.

Conflicts of interest

None

Ethical standards

The Birth to Twenty Study was granted approval by the Committee for Research on Human Subjects at the University of the Witwatersrand, Johannesburg.

References

1. de Onis M, Blössner M, Borghi E. Prevalence and trends of stunting among pre-school children, 1990–2020. *Public Health Nutr.* 2012; 15, 142-148.
2. Dewey KG, Begum K. Long-term consequences of stunting in early life. *Matern Child Nutr.* 2011; 7, 5-18.
3. Walker SP, Chang SM, Powell CA, Grantham-McGregor SM. Effects of early childhood psychosocial stimulation and nutritional supplementation on cognition and education in growth-stunted Jamaican children: prospective cohort study. *Lancet.* 2005; 366, 1804-7.
4. Victora CG, Adair L, Fall C, et al. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet.* 2008; 371, 340-57.
5. Victora CG, de Onis M, Hallal PC, Blossner M, Shrimpton R. Worldwide timing of growth faltering: revisiting implications for interventions. *Pediatrics* 2010; 125, e473-e480.
6. Cameron N, Preece MA, Cole TJ. Catch-up growth or regression to the mean? Recovery from stunting revisited. *Am J Hum Biol.* 2005; 17, 412-417.
7. Martorell R, Horta BL, Adair LS, et al. Weight gain in the first two years of life is an important predictor of schooling outcomes in pooled analyses from five birth cohorts from low- and middle-income countries. *J Nutr.* 2010; 140, 348-54.
8. Prentice AM, Ward KA, Goldberg GR, et al. Critical windows for nutritional interventions against stunting. *Am J Clin Nutr.* 2013; 97, 911-918.
9. Stein AD, Wang M, Martorell R, et al. Growth patterns in early childhood and final attained stature: Data from five birth cohorts from low- and middle-income countries. *Am J Hum Biol.* 2010; 22, 353-359.

10. Golden MH. Is complete catch-up possible for stunted malnourished children? *Eur J Clin Nutr.* 1994; 48 Suppl 1, S58-71.
11. Nelson CA, Thomas KM, de Haan M. *Neuroscience of cognitive development: the role of experience and the developing brain.* 2012; Hoboken, New Jersey: John Wiley & Sons.
12. Morgan B, Gibson KR. Nutritional and environmental interactions in brain development. In *Brain Maturation and Cognitive Development* (eds. Gibson KR, Peterson AC), 1991; pp. 91-106. Aldine de Bruyter, New York.
13. Shonkoff JP, Garner AS, Siegel BS, et al. The lifelong effects of early childhood adversity and toxic stress. *Pediatrics.* 2012; 129, e232-e246.
14. Glewwe P, King EM. The impact of early childhood nutritional status on cognitive development: does the timing of malnutrition matter? *World Bank Economic Review.* 2001; 15, 81-113.
15. Hoddinott J, Maluccio JA, Behrman JR, Flores R, Martorell R. Effect of a nutrition intervention during early childhood on economic productivity in Guatemalan adults. *Lancet.* 2008; 371, 411-6.
16. Mendez MA, Adair LS. Severity and timing of stunting in the first two years of life affect performance on cognitive tests in late childhood. *J Nutr.* 1999; 129, 1555-62.
17. Crookston BT, Schott W, Cueto S, et al. Postinfancy growth, schooling, and cognitive achievement: Young Lives. *Am J Clin Nutr.* 2013; 98, 1555-63.
18. Crookston BT, Penny ME, Alder SC, et al. Children who recover from early stunting and children who are not stunted demonstrate similar levels of cognition. *J Nutr.* 2010; 140, 1996-2001.

19. Crookston BT, Dearden KA, Alder SC, et al. Impact of early and concurrent stunting on cognition. *Matern Child Nutr.* 2011; 7, 397-409.
20. Casale D, Desmond C, Richter L. The association between stunting and psychosocial development among preschool children: a study using the South African Birth to Twenty cohort data. *Child Care Health Dev.* 2014; 40, 900-910.
21. Norris SA, Richter LM, Fleetwood SA. Panel studies in developing countries: case analysis of sample attrition over the past 16 years within the birth to twenty cohort in Johannesburg, South Africa. *Journal of International Development.* 2007; 19, 1143-1150.
22. Richter L, Norris S, Pettifor J, Yach D, Cameron N. Cohort Profile: Mandela's children: the 1990 Birth to Twenty study in South Africa. *Int J Epidemiol.* 2007; 36, 504-11.
23. Richter LM, Yach D, Cameron N, Griesel RD, Wet T. Enrolment into Birth to Ten (BTT): population and sample characteristics. *Paediatr Perinat Epidemiol.* 1995; 9, 109-120.
24. Frankenburg WK, Fandal AW, Thornton SM. Revision of Denver Prescreening Developmental Questionnaire. *J Pediatr.* 1987; 110, 653-657.
25. Hsiao C, Richter LM. Early mental development as a predictor of preschool cognitive and behavioral development in South Africa: the moderating role of maternal education in the Birth to Twenty Cohort. *Infants Young Child.* 2014; 27, 74-87.
26. Luiz DM, Foxcroft CD, Tukululu AN. The Denver II Scales and the Griffiths Scales of Mental Development: a correlational study. *J Child Adolesc Ment Health.* 2004; 16, 77-81.

27. WHO. *WHO Child Growth Standards: Methods and development: Head circumference-for-age, arm circumference-for-age, triceps skinfold-for-age and subscapular skinfold-for-age*. 2007; Geneva: World Health Organization.
28. Shisana O, Labadarios D, Rehle T, et al. *South African National Health and Nutrition Examination Survey (SANHANES-1)*. 2013; Cape Town: HSRC Press.
29. WHO. *Global Database on Child Growth and Malnutrition*. 2012; Available from: <http://www.who.int/nutgrowthdb/en/>.
30. Adair LS. Filipino children exhibit catch-up growth from age 2 to 12 years. *J Nutr*. 1999; 129, 1140-1148.
31. Berkman DS, Lescano AG, Gilman RH, Lopez SL, Black MM. Effects of stunting, diarrhoeal disease, and parasitic infection during infancy on cognition in late childhood: a follow-up study. *Lancet*. 2002; 359, 564-571.
32. Adair LS, Fall CH, Osmond C, et al. Associations of linear growth and relative weight gain during early life with adult health and human capital in countries of low and middle income: findings from five birth cohort studies. *Lancet*. 2013; 382, 525-34.
33. Boersma B, Wit JM. Catch-up growth. *Endocr Rev*. 1997; 18, 646-661.
34. Leroy JL, Ruel M, Habicht JP. Critical windows for nutritional interventions against stunting. *Am J Clin Nutr*. 2013; 98, 854-5.
35. Lundeen EA, Stein AD, Adair LS, et al. Height-for-age z scores increase despite increasing height deficits among children in 5 developing countries. *Am J Clin Nutr*. 2014; 100, 821-5.

TABLES

Table 1. Individual items comprising the R-DPDQ score, Bt20, age 5

ITEM
Asked of caregiver*
<ol style="list-style-type: none"> 1. Can X dress him/herself without help? 2. Can X play any simple board or card games? 3. Can X brush his/her teeth without help or supervision sometimes? 4. Can X get him/herself a bowl of cereal, spoon, dishing it out without making too much mess, and pouring milk (or other liquid) on it?
Interviewer tests⁺
<ol style="list-style-type: none"> 5. Build tower of blocks 6. Count blocks 1 7. Count blocks 5 8. Imitate vertical line 9. Copy a circle 10. Copy a cross 11. Copy a square – demonstrated 12. Pick a longer line 13. Draw a person - 3 parts 14. Draw a person – 6 parts 15. Knows use of objects – 3 16. Knows actions – 4 17. Understands prepositions – 4 18. Names colours – 1 19. Names colours – 4 20. Defines words – 5 21. Defines words – 7 22. Knows adjectives – 3 23. Opposites 24. Thumb wiggle 25. Balance on each foot 2 26. Balance on each foot 3 27. Balance on each foot 4 28. Balance on each foot 5 29. Balance on each foot 6 30. Hopping on one foot 31. Heel-to-toe walk 32. Interviewer rating of child’s speech (all or half understandable)

Notes:

*Responses: Yes (1), No (0), or No opportunity (0)

⁺Responses: Correct/Yes (1) or Incorrect/No (0)

Table 2. Mean values of the regression variables

	Mean (SD)/%	N
R-DPDQ 5y (score)	43.91 (4.70)	1019
Stunted 2y	20.1	1019
Stunted 5y	6.8	1019
HAZ 2y	-1.153 (1.095)	1019
HAZ 5y	-0.586 (0.949)	1019
Birth		
Female (%)	50.6	1019
Birth weight (g)	3071.8 (496.8)	1017
Socio-economic status		
Black African (%)	83.4	1019
Asset index 2y	3.86 (1.42)	856
Mother's age (years)	25.46 (6.24)	1019
Mother's schooling (years)	9.83 (2.55)	970
Home environment/caregiver inputs		
Mother is main caregiver 2y (%)	59.3	831
Birth order	2.03 (1.06)	1019
Child born within 24 months (%)	5.8	850
Caregiver plays 2y (%):		
- no time	4.0	855
- for less than an hr/day	36.7	855
- for more than an hr/day	59.3	855
Caregiver teaching child 2y (%)	78.1	850
Father(figure) plays 2y (%):		
- almost never	14.5	835
- once a week	21.8	835
- 2- 4 times/week	10.8	835
- every day	52.9	835
Child has toys (bought or made) 2y (%)	93.1	860
Change in SES 2y-4y		
Decrease in asset score	27.3	735
No change in asset score	47.3	735
Increase in asset score	25.3	735

Note: SD = standard deviation. *** p<0.01 ** p<0.05 * p<0.10.

Table 3. Stunting status and mean HAZ at 2 years and 5 years

	YEAR 2		YEAR 5	
	Prevalence (%)	HAZ	Prevalence (%)	HAZ
Not stunted	80.6	-0.751 (0.762)	Neither	79.3 -0.415 (0.767)
			Late incident	1.3 -2.216 (0.209)
Stunted	19.4	-2.72 (0.711)	Persistent	4.7 -2.464 (0.357)
			Catch up	14.7 -1.287 (0.503)
	100	n=1574	100	n=1574

Notes: Standard deviations in parentheses

Table 4. Estimates from cognitive scores regressions at 5y (OLS coefficients)

	I	II	III	IV	V
<u>Stunting status</u>					
Reference: Neither					
Persistent	-3.390*** (0.800)	-3.091*** (0.808)	-2.583*** (0.785)	-2.530*** (0.776)	-2.506*** (0.778)
Late incident	-0.357 (1.298)	-0.590 (1.292)	-0.754 (1.249)	-0.381 (1.247)	-0.416 (1.249)
Catch up	-2.258*** (0.432)	-1.978*** (0.438)	-1.739*** (0.425)	-1.602*** (0.423)	-1.607*** (0.425)
<u>Birth characteristics</u>					
Female		1.039*** (0.334)	0.975*** (0.323)	0.863*** (0.321)	0.867*** (0.322)
Birthweight		0.001* (0.000)	0.001* (0.000)	0.001* (0.000)	0.001* (0.000)
<u>Socio-economic status 2y</u>					
African			-1.589*** (0.529)	-1.798*** (0.546)	-1.783*** (0.549)
Asset score			0.495*** (0.119)	0.440*** (0.121)	0.476*** (0.136)
Mother's age			-0.045* (0.026)	-0.028 (0.037)	-0.029 (0.037)
Mother's schooling (yrs)			0.175*** (0.067)	0.110 (0.069)	0.107 (0.069)
<u>Home environment/caregiver 2y</u>					
Mother main caregiver				0.702** (0.321)	0.699** (0.321)
Birth order				-0.216 (0.231)	-0.206 (0.232)
Birth spacing				-1.271* (0.709)	-1.254* (0.712)
Reference: caregiver doesn't play with child					
Plays at least an hour/day				1.662** (0.813)	1.690** (0.815)
Plays more than an hour/day				0.976 (0.795)	1.001 (0.797)
Caregiver teaching child				0.002 (0.387)	0.005 (0.387)
Reference: father(figure) never plays with child					
Father(figure) plays once/week				0.135 (0.526)	0.146 (0.527)
Father(figure) plays 2-4/week				0.573 (0.630)	0.582 (0.631)
Father(figure) plays everyday				0.573 (0.473)	0.556 (0.474)
Toys (bought or made)				2.060*** (0.650)	2.052*** (0.651)
<u>Change in SES 2y-4y</u>					

Reference: Decrease in asset score					
No change in asset score					0.257 (0.381)
Increase in asset score					0.370 (0.480)
Constant	44.709*** (0.190)	42.102*** (1.137)	41.146*** (1.574)	38.366*** (1.759)	38.030*** (1.815)
N	666	666	666	666	666

Notes: Standard errors in parentheses. *** p<0.01 ** p<0.05 * p<0.10. The regressions are run on the sample of children who had non-missing data on the full set of explanatory variables in Regression IV.

Table 5: Mean characteristics for varying samples

	Sample of children with data on HAZ at 2y & 5y	Regression sample without covariates	Regression sample with covariates
R-DPDQ 5y (score)	-	43.91 (4.70)	44.14 (4.38)
HAZ 2y	-1.135 (1.084)	-1.153 (1.095)	-1.220 (1.105)
HAZ 5y	-0.664 (0.895)	-0.586 (0.949)	-0.570 (0.932)
Stunted 2y (%)	19.44	20.12	22.52
Stunted 5y (%)	6.04	6.77	6.16
Birth weight (g)	3072.30 (508.95)	3071.76 (496.82)	3084.06 (489.37)
Black (%)	75.60	83.42	89.34
Female (%)	52.35	50.64	49.85
Mother's age (yrs)	25.59 (6.19)	25.46 (6.24)	25.37 (6.35)
N	1574	1019	666

Note: Standard deviations in parentheses

Table 6. Estimates from unadjusted regressions, OLS coefficients (dependent variable = cognitive function scores at 5y)

	Full sample	Final regression sample
<u>Stunting status</u>		
Reference: Neither		
Persistent	-2.276*** (0.668)	-3.390*** (0.800)
Late incident	-0.563 (1.103)	-0.357 (1.298)
Catch up	-2.192*** (0.407)	-2.258*** (0.432)
N	1019	666

Notes: Standard errors in parentheses. *** p<0.01 ** p<0.05 * p<0.10.